arXiv:1602.08970v1 [hep-ex] 29 Feb 2016

Status and prospects for BSM ((N)MSSM) Higgs searches at the LHC

M.P. Casado[1] on behalf of the ATLAS and CMS collaborations.

[1]*Departamet de Física, Universitat Autònoma de Barcelona, Institut de Física d'Altes Energies and Barcelona Institute of Science and Technology, Spain.*

Talk presented at the International Workshop on Future Linear Colliders (LCWS15), Whistler, Canada, 2-6 November 2015

Abstract

Searches for Beyond the Standard Model Higgs processes in the context of Minimal Supersymmetric Standard Model and Next to MSSM are presented. The results are based on the first LHC run of pp collision data recorded by the ATLAS and CMS experiments at the CERN Large Hadron Collider at centre-of-mass energies of 7 and 8 TeV, corresponding to integrated luminosities of about 5 and 20 fb^{-1} respectively. Current searches constrain large parts of the parameter space. No evidence for BSM Higgs is found.

Keywords: BSM Higgs, Higgs, ATLAS, CMS, LHC

1. Introduction

The discovery of a new particle [1, 2] in the search for the Standard Model (SM) [3] Higgs boson at the Large Hadron Collider (LHC) [4], performed with the ATLAS [5] and CMS [6] detectors in July 2012, was an important achievement in the understanding of the electroweak symmetry breaking mechanism [7].

With this boson the Standard Model (SM) of particle physics is complete. However many effects remain unexplained in the SM, as the hierarchy problem or the existence of Dark Matter.

Moreover, there is no theoretical reason to restrict the model to only one Higgs boson: (1) the generation of fermion masses could also be realized by more bosons [8], (2) many theories include extra Higgs boson(s), as Supersymmetry (SUSY), models with axions, baryogenesis, neutrino masses, etc.

In this paper we will concentrate on Higgs bosons within the context of Minimal Supersymmetric Standard Model (MSSM) [9] and Next to Minimal Supersymmetric Standard Model (NMSSM) [10].



Figure 1: Invariant mass of the $\tau\tau$ pair in the electron, muon channel, no b-tag category in the $h/H/A \rightarrow \tau\tau$ analysis [12]. The main backgrounds are $Z \rightarrow \tau\tau$ processes and QCD events. There is good agreement between the observed data and the prediction.


ATLAS √s=8 TeV, ∫ L dt = 19.5 - 20.3 fb⁻¹
gluon fusion
φ → ττ
Obs 95% CL limit
Exp 95% CL limit
1 σ
2 σ
σ × BR(φ → ττ) [pb]
m_φ [GeV]
ATLAS √s=8 TeV, ∫ L dt = 19.5 - 20.3 fb⁻¹
b-associated production
φ → ττ
Obs 95% CL limit
Exp 95% CL limit
1 σ
2 σ
σ × BR(φ → ττ) [pb]
m_φ [GeV]

Figure 2: Observed and expected 95% CL limits on the cross-section times de branching ratio to a $\tau\tau$ pair as a function of the Higgs boson mass (gluon-fusion production on the left, b-associated production on the right) for the $h/H/A \to \tau\tau$ analysis [11]. No excess of events is found.

2. Searches within the Minimal Supersymmetric Standard Model

2.1. MSSM Neutral Higgs bosons at LHC

This search is important at high $\tan\beta$ (ratio of vacuum expectation values of the two Higgs doubles in the MSSM). The neutral Higgs field can be produced via gluon-fusion and in association with a b quark. The two most relevant final states are $h/H/A \to \tau\tau$ and bb. The $\tau\tau$ modes tend to be more sensitive due to the better performance of the tau-jet selection.

In the $h/H/A \to \tau\tau$ search a categorization is applied depending on the event properties ($\tau\tau$ decay and "b-tag"/"b-veto"). ATLAS performed the search using 8 TeV run 1 data [11] while CMS combined 7 and 8 TeV datasets [12].

Figure 1 presents the invariant mass of the $\tau\tau$ pair for the case in which one τ decays to electron and the other decays to a muon ("electron-muon" channel) and for the b-veto case in the CMS analysis. The main backgrounds are $Z \to \tau\tau$ processes and QCD events. There is good agreement between the observed data and the prediction. No excess of events is found.

Figure 2 shows the observed and expected 95% CL limits on the cross-section times branching ratio to a $\tau\tau$ pair as a function of the Higgs boson mass as obtained by ATLAS.

2.2. ATLAS and CMS search for $H^{\pm}$

This search covers the whole spectrum of $\tan\beta$. The $H^{\pm}$ can decay to $\tau\nu$, tb, cs, hW, etc. with different branching ratios depending on $\tan\beta$. The decay to $H^{\pm} \to \tau\nu$ is relevant in a large parameter range, specially for low $m_{H^{\pm}}$ (below m_{top}). For $m_{H^{\pm}}$ above m_{top} $H^{\pm} \to tb$ is the predominant decay.

ATLAS and CMS perform this search [13], [14] using the full 8 TeV Run 1 sample. The strategies are similar in both analyses. High and low mass $H^{\pm}$ categories are treated separately and a tau + missing E_T trigger is used to trigger events. The most important discriminating variable is presented in Fig. 3 for the ATLAS search. The 95% CL exclusion limits on $\tan\beta$ as a function of $m_H^{\pm}$ is shown for $m_H^{\pm} < m_{top}$ in Fig.4.

CMS has combined $H^{\pm} \to \tau\nu$ and $H^{\pm} \to tb$ channels. The observed and excluded limits on the $\tan\beta$-$m_{H^{\pm}}$ plane are presented in Fig.5. In this plot the excluded area by the discovery of the SM 125-GeV Higgs boson is drawn as a dotted red line and as a continuous red line if an error of $\pm 3GeV$ is included.

CMS has a recent analysis [15] searching for $m_H^{\pm} \to cs$. No excess of events is found.

2.3. Search for $A \to Zh$ at the LHC

This search is important at low $\tan\beta$ and has been performed both by ATLAS [16] and CMS [17]. Possible final states are $ll\tau\tau/llbb/\nu\nu bb$. It takes advantage of the $Z \to ll/Z \to \nu\nu/h$ resonances and ratios of the Higgs boson decays ($b\bar{b}/\tau\tau$).

Figure 6 shows the observed and expected 95% CL limits in the $\tan\beta$ - $\cos(\beta - \alpha)$ plane for two MSSM scenarios (Type I and Type II). The angle α is the CP-even Higgs mixing angle in the MSSM. No excess of events is observed.



$$m_T = \sqrt{2 p_T(\tau)\, E_T^{miss} (1 - \cos\Delta\phi(E_T^{miss}, \tau))}$$

Figure 3: Transverse invariant mass between visible τ decay products and the missing E_T of the event for the search of $H^{\pm} \to \tau\nu$ [13].



Figure 4: 95%CL exclusion limits on $\tan\beta$ as a function of $m_H^{\pm}$ is shown for $m_H^{\pm} < m_{top}$ in the $H^{\pm} \to \tau\nu$ search [13]. The interpretation is performed in the context of the m_h^{max} benchmark scenario [9] of the MSSM.



Figure 5: Observed and excluded limits on the $\tan\beta$-$m_H^{\pm}$ plane in the $H^{\pm} \to \tau\nu$ search [14] using the m_h^{mod-} benchmark scenario [9] of the MSSM. . The excluded area by the discovery of the SM 125-GeV Higgs boson is drawn as the red line.

2.4. Search for hh processes

This search is done by ATLAS [18] and CMS [19] in resonant and non-resonant Higgs boson pair production. The considered final states are $bb\gamma\gamma/bbbb/bb\tau\tau/WW\gamma\gamma$.

Figure 7 shows the non-resonant background fit in the $m_{\gamma\gamma}$ for one of the final states for the resonance mass hypothesis of 270 GeV in the CMS analysis.

2.5. Search for $H \to WW/ZZ$ processes

This search is done by ATLAS [20] and CMS [21].

The invariant masses obtained in the analysis allow to set upper 95% CL limits on the production cross-section. The different results from the different final states and the combination are presented in Fig. 8 for CMS.

3. Searches within the Next to Minimal Supersymmetric Standard Model

3.1. Search for $a \to \mu\mu$

This is a low mass search performed by CMS in [23]. The a boson is produced in gluon-fusion and searched for in the decay $\mu\mu$. In Fig. 9 the invariant mass of the dimuon pair is shown for barrel and endcap. The effect of a possible signal at 7 and 12 GeV is shown with the blue line. No excess of events is found in this search.



(a) Type-I 2HDM, $m_A = 300$ GeV



(b) Type-II 2HDM, $m_A = 300$ GeV

Figure 6: Observed and expected 95% CL limits in the $\tan\beta$ - $\cos(\beta-\alpha)$ plane for two MSSM scenarios [9] (Type I and Type II) in the $A \to Zh$ search [16]. The angle α is the CP-even Higgs mixing angle in the MSSM.



Figure 7: Non-resonant background fits in the $m_{\gamma\gamma}$ for one of the categories for the resonance mass hypothesis of 270 GeV in the *hh* CMS analysis [19].


CMS up to 5.1 fb-1 (7 TeV) + up to 19.7 fb-1 (8 TeV)
Combined
Combined (exp. ± 2σ)
H → WW → 2ℓ2ν
H → WW → ℓν qq
H → ZZ → 2ℓ2ν
H → ZZ → 4ℓ
H → ZZ → 2ℓ2τ
H → ZZ → 2ℓ2q
95% CL limit on σ/σSM
mH [GeV]

Figure 8: 95% CL on the ratio of cross-section over the SM cross-section for each of the contributing final states in the $H \to WW/ZZ$ search [21]. The theoretical cross section, σ_{SM}, is computed in [22].



Figure 9: Invariant mass of the dimuon pair for barrel (up) and endcap (bottom) in the $a \to \mu\mu$ search [23]. The effect of a possible signal at 7 and 12 GeV is shown with the blue line.

3.2. Search for $h \to aa \to \mu\mu\tau\tau/\mu\mu\mu\mu$

This search is performed by ATLAS [24] and CMS in [25]. The final states can be $\mu\mu\tau\tau/\mu\mu\mu\mu$. Fig. 10 shows the invariant mass of the dimuon pair for data and MC. On the top part of the Figure the light resonances are depicted showing the good understanding of the data.



Figure 10: Invariant mass of the dimuon pair for data and MC in the $h \to aa \to \mu\mu\tau\tau/\mu\mu\mu\mu$ search [24]. On the top part of the Figure the light resonances are depicted showing the good understanding of the data.

4. Conclusions

Current searches constrain large parts of the parameter space. So far there is no evidence for Beyond the Standard Model Higgs. However, there are still many possibilities to explore and many searches are still starting up, and this will be a relevant area in Run 2.

References

[1] ATLAS Collaboration, *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, Phys. Lett. **B 716** (2012) 1, arXiv:1207.7214

[2] CMS Collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, Phys. Lett. **B 716** (2012) 30, arXiv:1207.7235

[3] S. L. Glashow, *Partial-symmetries of weak interactions* , Nucl. Phys. 22 (1961) 579
S. Weinberg, *A model of leptons*, Phys. Rev. Lett. 19 (1967) 1264
A. Salam, *Elementary Particle Theory* , Almqvist and Wiksells (Stockholm), (1968) 367

[4] L. Evans, P. Bryant (Eds.), *LHC Machine*, JINST **3** S08001 (2008)

[5] ATLAS Collaboration, *The ATLAS Experiment at the CERN Large Hadron Collider*, JINST **3** S08003 (2008)

[6] CMS Collaboration, *The CMS Experiment at the CERN LHC*, JINST **3** S08004 (2008).

[7] F. Englert and R. Brout, *Broken symmetry and the mass of gauge vector mesons*, Phys. Rev. Lett. **13** (1964) 321
P. W. Higgs, *Broken symmetries, massless particles and gauge fields*, Phys. Lett. **12** (1964) 132
P. W. Higgs, *Broken symmetries and the masses of gauge bosons*, Phys. Rev. Lett. **13** (1964) 508
G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, *Global conservation laws and massless particles*, Phys. Rev. Lett. **13** (1964) 585
P. W. Higgs, *Spontaneous symmetry breakdown without massless bosons*, Phys. Rev. **145** (1966) 1156
T. W. B. Kibble, *Symmetry breaking in non-Abelian gauge theories*, Phys. Rev. **155** (1967) 1554

[8] D. Ghosh, R. Sandeepan Gupta, G. Perez *Is the Higgs Mechanism of Fermion Mass Generation a Fact? A Yukawa-less First-Two-Generation Model*, arXiv: 1508.01501

[9] M. S. Carena, S. Heinemeyer, C. Wagner, and G. Weiglein, *Suggestions for improved benchmark scenarios for Higgs boson searches at LEP-2* , arXiv:hep-ph/9912223
M. Carena et al., *MSSM Higgs Boson Searches at the LHC: Benchmark Scenarios after the Discovery of a Higgs-like Particle*, Eur. Phys. J. C 73 (2013) 2552, arXiv:1302.7033

[10] M. Maniatis, *The Next-to-Minimal Supersymmetric extension of the Standard Model reviewed*, Int. J. Mod. Phys. A 25 (2010) 3505, arXiv:0906.0777

[11] ATLAS Collaboration, *Search for neutral Higgs bosons of the minimal supersymmetric standard model in pp collisions at* $\sqrt{s} = 8$ *TeV with the ATLAS detector*, JHEP11(2014) 056, arXiv: 1409.6064

[12] CMS Collaboration, *Search for neutral MSSM Higgs bosons decaying to a pair of tau leptons in pp collisions*, JHEP10 (2014) 160, arXiv: 1408.3316

[13] ATLAS Collaboration, *Search for charged Higgs bosons decaying via* $H^{\pm} \to \tau\nu$ *in fully hadronic final states using pp collision*

data at $\sqrt{s}$ *= 8 TeV with the ATLAS detector*, JHEP03 (2015) 088, arXiv: 1412.6663

[14] CMS Collaboration, *Search for a charged Higgs boson in pp collisions at* $\sqrt{s}$ *= 8 TeV*, arXiv: 1508.07774, accepted by JHEP

[15] CMS Collaboration, *Search for* H^+ *to cs-bar decay*, CMS-PAS-HIG-13-035

[16] ATLAS Collaboration, *Search for a CP-odd Higgs boson decaying to Zh in pp collisions at* $\sqrt{s}$ *=* $8TeV$ *with the ATLAS detector*, Phys. Lett. **B 744** (2015), 163-183

[17] CMS Collaboration, *Search for a pseudoscalar boson decaying into a Z boson and the 125 GeV Higgs boson in* $l^+l^-b\bar{b}$ *final states*, Phys. Lett. **B 748** (2015), 221-243, arXiv: 1504.04710

[18] ATLAS Collaboration, *Searches for Higgs boson pair production in the* $hh \rightarrow bb\tau\tau, \gamma\gamma WW*, \gamma\gamma bb, bbbb$ *channels with the ATLAS detector*, Phys. Rev. **D 92**, 092004 (2015), arXiv: 1509.04670

[19] CMS Collaboration, *Search for resonant HH production in 2gamma+2b channel*, CMS-PAS-HIG-13-032, correction in https://twiki.cern.ch/twiki/bin/view/CMSPublic/Hig13032TWiki

[20] ATLAS Collaboration, *Search for an additional, heavy Higgs boson in the* $H \rightarrow ZZ$ *decay channel at* $\sqrt{s}$ *= 8 TeV in pp collision data with the ATLAS detector*, Eur. Phys. J. **C 76** (2016) 45, arXiv: 1507.05930

[21] CMS Collaboration, *Search for a Higgs boson in the mass range from 145 to 1000 GeV decaying to a pair of W or Z bosons*, JHEP10 (2015) 144, arXiv: 1504.00936

[22] LHC Higgs Cross Section Working Group, *Handbook of LHC Higgs cross sections: 3. Higgs Properties*, arXiv: 1307.1347

[23] CMS Collaboration, *Search for a light pseudoscalar Higgs boson in the dimuon decay channel in pp collisions at* $\sqrt{s}$ *= 7 TeV*, Phys. Rev. Lett. **109** (2012) 121801, arXiv: 1206.6326

[24] ATLAS Collaboration, *Search for Higgs bosons decaying to aa in the* $\mu\mu\tau\tau$ *final state in pp collisions at* $\sqrt{s}$ *= 8 TeV with the ATLAS experiment*, Phys. Rev. **D 92**, 052002 (2015), arXiv: 1505.01609

[25] CMS Collaboration, *A search for pair production of new light bosons decaying into muons*, Phys. Rev. Lett. **752** (2016) 146,arXiv: 1506.00424